

10. Kissane DW, Maj M, Sartorius N Depression and Cancer. Wiley Blackwell Oxford 2011; 300 s.
11. Maatouk I, He S, Becker N, et al. Association of resilience with health-related quality of life and depression in multiple myeloma and its precursors: results of a German cross-sectional study. *BMJ Open* 2018; 8: e021376.
12. Kieley F, Cran A, Finnerty D, et al. Self-Reported Quality of Life and Symptom Burden in Ambulatory Patients With Multiple Myeloma on Disease-Modifying Treatment. *Am J Hosp Palliat Care* 2017; 34: 671–676.
13. Lamers J, Hartmann M, Goldschmidt H, et al. Psychosocial support in patients with multiple myeloma at time of diagnosis: who wants what? *Psychooncology* 2013; 22: 2313–2320.
14. Oberoi D, White V, Seymour J, et al. The course of anxiety, depression and unmet needs in survivors of diffuse large B cell lymphoma and multiple myeloma in the early survivorship period. *J Cancer Surviv* 2017; 11: 329–338.
15. Mols F, Husson O, Roukema JA, et al. Depressive symptoms are a risk factor for all-cause mortality: results from a prospective population-based study among 3,080 cancer survivors from the PROFILES registry. *J Cancer Surviv* 2013; 7: 484–492.
16. Niazi S, Frank RD, Sharma M, et al. Impact of psychiatric comorbidities on health care utilization and cost of care in multiple myeloma. *Blood Adv* 2018; 2: 1120–1128.
17. El-Jawahri A, Chen YB, Brazauskas R, et al. Impact of pre-transplant depression on outcomes of allogeneic and autologous hematopoietic stem cell transplantation. *Cancer* 2017; 123: 1828–1838.
18. Svěrák T, Skřivanová K, Anderková L, et al. Screening of psychological distress 4,5 years after diagnosis in breast cancer patients compared to healthy population *Klinická onkologie* 2016; 29: 210–215.
19. Molassiotis A, Wilson B, Blair S, et al. Unmet supportive care needs, psychological well-being and quality of life in patients living with multiple myeloma and their partners. *Psychooncology* 2011; 20: 88–97.
20. Moghaddam N, Coxon H, Nabarro S, et al. Unmet care needs in people living with advanced cancer: a systematic review. *Support Care Cancer* 2016; 24: 3609–3622.
21. Bužgová R, Hajnová Fukasová E, Sikorová L et al. Association between unmet needs and quality of life in hospitalised cancer patients no longer receiving anti-cancer treatment. *Eur. J Cancer Care* 2014; 23: 685–694.
22. Látalová K, Praško J, Kamarádová D, et al. Posttraumatická stresová porucha. *Praktický lékař* 2014; 94: 172–176.
23. Mehnert A, Lehmann C, Graefen M, et al. Depression, anxiety, post-traumatic stress disorder and health-related quality of life and its association with social support in ambulatory prostate cancer patients. *Eur J Cancer Care (Engl)* 2010; 19: 736–745.
24. Thekdi SM, Milbury K, Spelman A, et al. Posttraumatic Stress and Depressive Symptoms in Renal Cell Carcinoma: Association with Quality of Life and Utility of Single Item Distress Screening. *Psychooncology* 2015; 24: 1477–1484.
25. Geffen DB, Blaustein A. Post-traumatic stress disorder and quality of life in long-term survivors of Hodgkin's disease and non-Hodgkin's lymphoma in Israel. *Leuk Lymphoma* 2003; 44: 1925–1929.
26. Smith SK, Zimmerman S, Williams CS, et al. Post-traumatic stress symptoms in long-term non-Hodgkin's lymphoma survivors: does time heal? *J Clin Oncol* 2011; 29: 4526–4533.
27. Rodin G, Yuen D, Mischitelle A, et al. Traumatic stress in acute leukemia. *Psychooncology* 2013; 22: 299–307.
28. Jawahri AR, Vandusen HB, Traeger LN, et al. Quality of life and mood predict posttraumatic stress disorder after hematopoietic stem cell transplantation. *Cancer* 2016; 122: 806–812.
29. Avery DH, Shah SH, Eder DN, et al. Nocturnal sweating and temperature in depression. *Acta Psychiatr Scand* 1999; 100: 295–301.
30. Rietsema WJ. Post-traumatic stress disorder as a cause of night sweats. *Am Fam Physician* 2003; 68: 806–808.
31. Mold JW, Holtzclaw BJ, McCarthy L. Night sweats: a systematic review of the literature. *J Am Board Fam Med* 2012; 25: 878–893.
32. Oka T. Psychogenic fever: how psychological stress affects body temperature in the clinical population. *Temperature (Austin)* 2015; 2: 368–378.
33. Takakazu O. Psychogenic fever: how psychological stress affects body temperature in the clinical population. *Temperature (Austin)* 2015; 2: 368–378.
34. Steen R, Dahl AA, Hess SL et al. A study of chronic fatigue in Norwegian cervical cancer survivors. *Gynecol Oncol* 2017; 146: 630–635.
35. Nebiker L, Lichtenstein E, Minghetti A, et al. Moderating Effects of Exercise Duration and Intensity in Neuromuscular vs. Endurance Exercise Interventions for the Treatment of Depression: A Meta-Analytical Review. *Front Psychiatry* 2018; 9: 305.
36. Rebar RA, Stanton R, Geard D, et al. A meta-analysis of the effect of physical activity on depression and anxiety in non-clinical adult populations. *Health Psychology Review* 2015; 9: 366–378.
37. Werneck AO, Oyeyemi AL, Silva Dr. Physical activity and depression: is 150 min/week of moderate to vigorous physical activity a necessary threshold for decreasing risk of depression in adults? Different views from the same data. *Soc Psychiatry Psychiatr Epidemiol* 2018; 53: 323–324.
38. Marker RJ, Cox-Martin E, Jankowski CM, et al. Evaluation of the effects of a clinically implemented exercise program on physical fitness, fatigue, and depression in cancer survivors. *Support Care Cancer* 2018; 26: 1861–1869.
39. Hadrabová M, Janíková A. Význam fyzické aktivity pro přežití a kvalitu života u pacientů s lymfoproliferativním onemocněním. *Transfuzie a Hematologie dnes* 2018; 3: 182–193.
40. Janíková A, Radvanský J, Vysoký R, et al. Význam fyzické aktivity u pacientů s hematologickými malignitami. *Transfuzie a hematologie dnes* 2012; 18: 31–38.