

10. Santema BT, Chan MMY, Tromp J et al. The influence of atrial fibrillation on the levels of NT-proBNP versus GDF-15 in patients with heart failure. *Clin Res Cardiol.* 2020; 109(3): 331–338.
11. Sharma A, Stevens SR, Lucas J et al. Utility of Growth Differentiation Factor-15, A Marker of Oxidative Stress and Inflammation, in Chronic Heart Failure: Insights From the HF-ACTION Study. *JACC Heart Fail.* 2017; 5(10): 724–734.
12. Bouabdallaoui N, Claggett B, Zile MR et al. Growth differentiation factor-15 is not modified by sacubitril/valsartan and is an independent marker of risk in patients with heart failure and reduced ejection fraction: the PARADIGM-HF trial. *Eur J Heart Fail.* 2018; 20(12): 1701–1709.
13. Cotter G, Voors AA, Prescott MF et al. Growth differentiation factor 15 (GDF-15) in patients admitted for acute heart failure: results from the RELAX-AHF study. *Eur J Heart Fail.* 2015; 17(11): 1133–1143.
14. Anand IS, Kempf T, Rector TS et al. Serial measurement of growth-differentiation factor-15 in heart failure: relation to disease severity and prognosis in the Valsartan Heart Failure Trial. *Circulation.* 2010; 122(14): 1387–1395.
15. Sunman H, Canpolat U, Yorgun H et al. Association between reverse electrical remodeling and cardiac fibrosis markers in patients with cardiac resynchronization therapy. *Turk Kardiyol Dern Ars.* 2018; 46(2): 84–91.
16. Foley PW, Stegemann B, Ng K et al. Growth differentiation factor-15 predicts mortality and morbidity after cardiac resynchronization therapy. *Eur Heart J.* 2009; 30(22): 2749–2757.
17. Hindricks G, Potpara T, Dagres N et al. 2020 ESC Guidelines for the diagnosis and management of atrial fibrillation developed in collaboration with the European Association of Cardio-Thoracic Surgery (EACTS). *Eur Heart J.* 2020.
18. Hijazi Z, Oldgren J, Lindback J et al. The novel biomarker-based ABC (age, biomarkers, clinical history)-bleeding risk score for patients with atrial fibrillation: a derivation and validation study. *Lancet.* 2016; 387(10035): 2302–2311.
19. Esteve-Pastor MA, Rivera-Caravaca JM, Roldan V et al. Long-term bleeding risk prediction in 'real world' patients with atrial fibrillation: Comparison of the HAS-BLED and ABC-Bleeding risk scores. The Murcia Atrial Fibrillation Project. *Thromb Haemost.* 2017; 117(10): 1848–1858.
20. Wallentin L, Lopes RD, Hanna M et al. Efficacy and safety of apixaban compared with warfarin at different levels of predicted international normalized ratio control for stroke prevention in atrial fibrillation. *Circulation.* 2013; 127(22): 2166–2176.
21. Connolly SJ, Ezekowitz MD, Yusuf S et al. Dabigatran versus warfarin in patients with atrial fibrillation. *N Engl J Med.* 2009; 361(12): 1139–1151.
22. Berg DD, Ruff CT, Jarolim P et al. Performance of the ABC Scores for Assessing the Risk of Stroke or Systemic Embolism and Bleeding in Patients With Atrial Fibrillation in ENGAGE AF-TIMI 48. *Circulation.* 2019; 139(6): 760–771.
23. Esteve-Pastor MA, Rivera-Caravaca JM, Shantsila A et al. Assessing Bleeding Risk in Atrial Fibrillation Patients: Comparing a Bleeding Risk Score Based Only on Modifiable Bleeding Risk Factors against the HAS-BLED Score. The AMADEUS Trial. *Thromb Haemost.* 2017; 117(12): 2261–2266.
24. Hijazi Z, Oldgren J, Lindback J et al. A biomarker-based risk score to predict death in patients with atrial fibrillation: the ABC (age, biomarkers, clinical history) death risk score. *Eur Heart J.* 2018; 39(6): 477–485.
25. Kempf T, Bjorklund E, Olofsson S et al. Growth-differentiation factor-15 improves risk stratification in ST-segment elevation myocardial infarction. *Eur Heart J.* 2007; 28(23): 2858–2865.
26. Wallentin L, Lindholm D, Siegbahn A et al. Biomarkers in relation to the effects of ticagrelor in comparison with clopidogrel in non-ST-elevation acute coronary syndrome patients managed with or without in-hospital revascularization: a substudy from the Prospective Randomized Platelet Inhibition and Patient Outcomes (PLATO) trial. *Circulation.* 2014; 129(3): 293–303.
27. Wallentin L, Lindhagen L, Arnstrom E et al. Early invasive versus non-invasive treatment in patients with non-ST-elevation acute coronary syndrome (FRISC-II): 15 year follow-up of a prospective, randomised, multicentre study. *Lancet.* 2016; 388(10054): 1903–1911.
28. Wollert KC, Kempf T, Peter T et al. Prognostic value of growth-differentiation factor-15 in patients with non-ST-elevation acute coronary syndrome. *Circulation.* 2007; 115(8): 962–971.
29. Widera C, Pencina MJ, Bobadilla M et al. Incremental prognostic value of biomarkers beyond the GRACE (Global Registry of Acute Coronary Events) score and high-sensitivity cardiac troponin T in non-ST-elevation acute coronary syndrome. *Clin Chem.* 2013; 59(10): 1497–1505.
30. Morrow DA, Scirica BM, Karwatowska-Prokopczuk E et al. Effects of ranolazine on recurrent cardiovascular events in patients with non-ST-elevation acute coronary syndromes: the MERLIN-TIMI 36 randomized trial. *JAMA.* 2007; 297(16): 1775–1783.
31. Collet JP, Thiele H, Barbato E et al. 2020 ESC Guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation. *Eur Heart J.* 2020.
32. Kahli A, Guenancia C, Zeller M et al. Growth differentiation factor-15 (GDF-15) levels are associated with cardiac and renal injury in patients undergoing coronary artery bypass grafting with cardiopulmonary bypass. *PLoS One.* 2014; 9(8): e105759.
33. Munk PS, Valborgland T, Butt N, Larsen AI. Response of growth differentiation factor-15 to percutaneous coronary intervention and regular exercise training. *Scand Cardiovasc J.* 2011; 45(1): 27–32.

Rádi vám pomáhamé v on-line vzdělávání

23

on-line kurzů a kongresů

160

přednášek

1340

vydaných certifikátů

7200

účastníků

