

25. Gierlotka M, Tajstra M, Gašior M et al. Impact of chronic total occlusion artery on 12-month mortality in patients with non-ST-segment elevation myocardial infarction treated by percutaneous coronary intervention (from the PL-ACS Registry). *Int J Cardiol.* 2013; 168(1): 250–254. doi:10.1016/j.ijcard.2012.09.086.
26. Watanabe H, Morimoto T, Shiomi H et al. Chronic total occlusion in a non-infarct-related artery is closely associated with increased five-year mortality in patients with ST-segment elevation acute myocardial infarction undergoing primary percutaneous coronary intervention (from the CREDO-Kyoto AML registry). *EuroIntervention.* 2017; 12(15): e1874–e1882. doi:10.4244/EIJ-D-15–00421.
27. Fujii T, Sakai K, Nakano M et al. Impact of the origin of the collateral feeding donor artery on short-term mortality in ST-elevation myocardial infarction with comorbid chronic total occlusion. *Int J Cardiol.* 2016; 218: 158–163. doi:10.1016/j.ijcard.2016.05.023.
28. Henriques JP, Hoehbers LP, Råmunddal T et al. Percutaneous Intervention for Concurrent Chronic Total Occlusions in Patients With STEMI: The EXPLORE Trial. *J Am Coll Cardiol.* 2016; 68(15): 1622–1632. doi:10.1016/j.jacc.2016.07.744.
29. Hachamovitch R, Hayes SW, Friedman JD et al. Comparison of the short-term survival benefit associated with revascularization compared with medical therapy in patients with no prior coronary artery disease undergoing stress myocardial perfusion single photon emission computed tomography. *Circulation.* 2003; 107(23): 2900–2907. doi:10.1161/01.CIR.0000072790.23090.41.
30. Galassi AR, Brilakis ES, Boukhris M et al. Appropriateness of percutaneous revascularization of coronary chronic total occlusions: an overview. *Eur Heart J.* 2016;37(35):2692–2700. doi:10.1093/eurheartj/ehv391.
31. Galassi AR, Werner GS, Boukhris M et al. Percutaneous recanalisation of chronic total occlusions: 2019 consensus document from the EuroCTO Club. *EuroIntervention.* 2019; 15(2): 198–208. doi:10.4244/EIJ-D-18–00826.
32. Schumacher SP, Kockx M, Stuijzand WJ et al. Ischaemic burden and changes in absolute myocardial perfusion after chronic total occlusion percutaneous coronary intervention. *EuroIntervention.* 2020; 16(6): e462–e471. Published 2020 Aug 7. doi:10.4244/EIJ-D-19–00631.
33. Galassi AR, Werner GS, Tomasello SD et al. Prognostic value of exercise myocardial scintigraphy in patients with coronary chronic total occlusions. *J Interv Cardiol.* 2010; 23(2): 139–148. doi:10.1111/j.1540–8183.2010.00527.
34. Löffler AI, Kramer CM. Myocardial Viability Testing to Guide Coronary Revascularization. *Interv Cardiol Clin.* 2018; 7(3): 355–365. doi:10.1016/j.iccl.2018.03.005.
35. Bucciarelli-Ducci C, Auger D, Di Mario C et al. CMR Guidance for Recanalization of Coronary Chronic Total Occlusion. *JACC Cardiovasc Imaging.* 2016; 9(5): 547–556. doi:10.1016/j.jcmg.2015.10.025.
36. Gerbay A, Youssef E, Vola M et al. Prediction of recovery after revascularization in chronic Coronary Total Occlusion (CTO) patients. Adenosine or low-dose dobutamine stress with LGE CMR: which is the best combination? *J Cardiovasc Magn Reson.* 2015; 17(Suppl 1): P114. Published 2015 Feb 3. doi:10.1186/1532–429X-17-S1-P114.
37. Pica S, Di Giovine G, Bollati M et al. Cardiac magnetic resonance for ischaemia and viability detection. Guiding patient selection to revascularization in coronary chronic total occlusions: The CARISMA\_CTO study design. *Int J Cardiol.* 2018; 272: 356–362. doi:10.1016/j.ijcard.2018.08.061.
38. Okamura A, Yamane M, Muto M et al. Complications during retrograde approach for chronic coronary total occlusion: Sub-analysis of Japanese multicenter registry. *Catheter Cardiovasc Interv.* 2016; 88(1): 7–14. doi:10.1002/ccd.26317.
39. Galassi AR, Sianos G, Werner GS et al. Retrograde Recanalization of Chronic Total Occlusions in Europe: Procedural, In-Hospital, and Long-Term Outcomes From the Multicenter ERCTO Registry. *J Am Coll Cardiol.* 2015; 65(22): 2388–2400. doi:10.1016/j.jacc.2015.03.566.
40. Mashayekhi K, Behnes M, Akin I, Kaiser T, Neuser H. Novel retrograde approach for percutaneous treatment of chronic total occlusions of the right coronary artery using ipsilateral collateral connections: a European centre experience. *EuroIntervention.* 2016;11(11):e1231–e1236. doi:10.4244/EIJV11111A244.
41. Brilakis ES, Grantham JA, Rinfret S et al. A percutaneous treatment algorithm for crossing coronary chronic total occlusions. *JACC Cardiovasc Interv.* 2012; 5(4): 367–379. doi:10.1016/j.jcin.2012.02.006.

# Recenzované články na téma **COVID-19**,

→  
které vycházejí v našich odborných časopisech,  
najdete v plném znění na  
**www.solen-covid.cz**

Akutní infekce horních cest dýchacích a jejich management / Akutní respirační infekce – symptomatická samoléčba u dospělých / Mohou sérologické metody pomoci při diagnostice covid-19 v případě opakované falešné negativity výsledku PCR testu? / Nejčastější chyby při používání osobních ochranných pomůcek / Koagulopatie asociovaná s onemocněním covid-19 / Etické konotace léčby onemocnění covid-19 / Molekula erdostein společnosti Recipharm byla s pozitivním výsledkem testována jako součást léčby pacientů s covid-19 / Antioxidační a bronchoprotektivní účinek erdosteinu / Suplementácia vitamínu D ako dôležitý faktor v prevencii a liečbe ochorenia covid-19: aké máme dôkazy? / Akutní infekce, žilní trombóza a doporučená trombo profylaxe / Zpráva z TCT 2020 / Pětileté zkušenosti z ECMO programu regionálního kardiocentra bez kardiochirurgie / Jsou vakcíny bezpečné? ... A ŘADA DALŠÍCH ČLÁNKŮ

