


3. Wilson ML, Weinstein MP, Reinier LG, et al. Controlled comparison of the BacT/Alert and BACTEC 660/730 nonradiometric blood culture systems. *J Clin Microbiol.* 1992;30:323-9.
4. Baron EJ, Scott JD, Tompkins LS. Prolonged incubation and extensive subculturing do not increase recovery of clinically significant microorganisms from standard automated blood cultures. *Clin Infect Dis.* 2005;41:1677-1680.
5. Petti CA, Bhally HS, Weinstein MP, et al. Utility of extended blood culture incubation for isolation of *Haemophilus*, *Actinobacillus*, *Cardiobacterium*, *Eikenella*, and *Kingella* organisms: a retrospective multicenter evaluation. *J Clin Microbiol.* 2006;44:257-259.
6. Kolar M, Htoutou Sedlakova M, Urbanek K, et al. Implementation of Antibiotic Stewardship in a University Hospital Setting. *Antibiotics.* 2021;10(1):93.
7. Riedel S, Bourbeau P, Swartz B, et al. Timing of specimen collection for blood cultures from febrile patients with bacteremia. *J Clin Microbiol.* 2008;46(4):1381-5.
8. Lamy B, Dargère S, Arendrup MC, et al. How to Optimize the Use of Blood Cultures for the Diagnosis of Bloodstream Infections? A State-of-the-Art. *Front Microbiol.* 2016;12(7):697.
9. Lee A, Mirrett S, Reller LB, Weinstein MP. Detection of Bloodstream Infections in Adults: How Many Blood Cultures Are Needed? *J Clin Microbiol.* 2007;45:3546-3548.
10. Li J, Plorde JJ, Carlson LG. Effects of volume and periodicity on blood cultures. *J Clin Microbiol.* 1994;32:2829-2831.
11. Scheer C, Idelevich EA, Annane D, et al. Reported time to blood culture results in hospitals with 24/7 microbiological service compared to hospitals with limited microbiological service: results from an international survey in 870 hospitals. *ECCMID 2023.*
12. Drevinek P, Hurych J, Antuskova M, et al. Direct detection of ESKAPEc pathogens from whole blood using the T2Bacteria Panel allows early antimicrobial stewardship intervention in patients with sepsis. *Microbiologyopen.* 2021;10(3):e1210.
13. Salar-Vidal L, Fernández-Roblas F, Gadea-Gironés I, et al. T2 magnetic resonance for selected adult patients: experience in a tertiary hospital with sepsis code. *ECCMID 2023.*
14. Cheng MP, Stenstrom R, Paquette K, et al. Blood culture results before and after antimicrobial administration in patients with severe manifestations of sepsis: A diagnostic study. *Ann of Intern Med.* 2019;171(8):547-554.
15. Singhal N, Kumar M, Kanaujia PK, et al. MALDI-TOF mass spectrometry: an emerging technology for microbial identification and diagnosis. *Front Microbiol.* 2015;6:791.
16. Torres-Sangiao E, Leal Rodríguez C, García-Riestra C. Application and Perspectives of MALDI-TOF Mass Spectrometry in Clinical Microbiology Laboratories. *Microorganisms.* 2021;9(7):1539.
17. Morgenthaler NG, Kostrzewa M. Rapid identification of pathogens in positive blood culture of patients with sepsis: review and meta-analysis of the performance of the sepsityper kit. *Int J Microbiol.* 2015;2015:827416.
18. Bazzi AM, Rabaan AA, El Edaily Z, et al. Comparison among four proposed direct blood culture microbial identification methods using MALDI-TOF MS. *J Infect Public Health.* 2017;10(3):308-15.
19. Azrad M, Keness Y, Nitzan O, et al. Cheap and rapid in-house method for direct identification of positive blood cultures by MALDI-TOF MS technology. *BMC Infect Dis.* 2019;19(1):72.
20. La Scola B, Raoult D. Direct Identification of Bacteria in Positive Blood Culture Bottles by Matrix-Assisted Laser Desorption Ionisation Time-of-Flight Mass Spectrometry. *PLoS One.* 2009;4(11):e8041.
21. Ferreira L, Sánchez-Juanes F, Muñoz-Bellido J et al. Rapid method for direct identification of bacteria in urine and blood culture samples by matrix-assisted laser desorption ionization time-of-flight mass spectrometry: intact cell vs. extraction method. *Clin Microbiol Inf.* 2011;17(7):1007-1012.
22. Juiz P, Almela M, Melción C, et al. A comparative study of two different methods of sample preparation for positive blood cultures for the rapid identification of bacteria using MALDI-TOF MS. *Eur J Clin Microbiol Infect Dis.* 2011;31(7):1353-1358.
23. Yonetani S, Ohnishi H, Ohkusu K, et al. Direct identification of microorganisms from positive blood cultures by MALDI-TOF MS using an in-house saponin method. *Int J Infect Dis.* 2016;52:37-42.
24. European Committee on Antimicrobial Susceptibility Testing. Methodology – EUCAST rapid antimicrobial susceptibility testing (RAST) directly from positive blood culture bottles. [cit. 2023-01-04] : Available from www: https://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/RAST/2023/2023_June/EUCAST_RAST_Breakpoint_Table_v_6.1_final_PDF.pdf
25. Kreidl P, Kirchner T, Fille M, et al. Antibiotic resistance of blood cultures in regional and tertiary hospital settings of Tyrol, Austria (2006-2015): Impacts & trends. *PLoS ONE.* 2019;14(10):e0223467.
26. Karlowsky JA, Jones ME, Draghi DC, et al. Prevalence and antimicrobial susceptibilities of bacteria isolated from blood cultures of hospitalized patients in the United States in 2002. *Ann Clin Microbiol Antimicrob.* 2004;10:3-7.
27. Wilson J, Elgohari S, Livermore DM, et al. Trends among pathogens reported as causing bacteraemia in England, 2004–2008. *Clin Microbiol Infect.* 2011;17:451-458.
28. Htoutou Sedláková M, Fišerová K, Kolář M. Původci bakteriemií ve Fakultní nemocnici Olomouc. *Klin mikrobiol inf lék.* 2020;26(1):4-11.
29. Banerjee R, Humphries R. Rapid Antimicrobial Susceptibility Testing Methods for Blood Cultures and Their Clinical Impact. *Front Med. (Lausanne)* 2021;10(8):635831.
30. Fiserova K, Htoutou Sedlakova M, Kolar M. The impact of rapid blood cultures identification by MALDI-TOF MS technology. *ECCMID 2023*, the accepted abstract (Poster number P1564).

Vnitřní lékařství

na rok 2024

Už máte předplaceno?

 SOLEN
let s vámi




**PŘEDPLATNÝM
ČASOPISU
NA ROK 2024
ZÍSKÁTE**

8 čísel ve vaší schránce

Tematická suplementa

**Přístup k archivu PDF
s praktickými tabulkami
pro internisty**



OBJEDNÁVEJTE

www.casopisvnitrnilekarstvi.cz

predplatne@solen.cz

585 204 335

