

LITERATURA

1. Baumgartner RN, Wayne SJ, Waters DL, et al. Sarcopenic obesity predicts instrumental activities of daily living disability in the elderly. *Obes Res.* 2004;12:1995-2004.
2. Mathus-Vliegen EM. Obesity Management Task Force of the European Association for the Study of Obesity. Prevalence, pathophysiology, health consequences and treatment options of obesity in the elderly: a guideline. *Obes Facts.* 2012;5(3):460-83
3. Prado CM, Batsis JA, Donini LM, et al. Sarcopenic obesity in older adults: a clinical overview. *Nat Rev Endocrinol.* 2024 May;20(5):261-277. doi: 10.1038/s41574-023-00943-z. Epub 2024 Feb 6. PMID: 38321142.
4. Poggiogalle E, Lubrano C, Sergi G, et al. Sarcopenic obesity and metabolic syndrome in adult Caucasian subjects. *J Nutr Health Aging.* 2016;20(9):958e63. <https://doi.org/10.1007/s12603-015-0638-1>.
5. Donini LM, Busetto L, Bischoff SC, et al. Definition and Diagnostic Criteria for Sarcopenic Obesity: ESPEN and EASO Consensus Statement. *Obes Facts.* 2022;15(3):321-335.
6. Donini LM, Busetto L, Bauer JM, et al. Critical appraisal of definitions and diagnostic criteria for sarcopenic obesity based on a systematic review. *Clin Nutr.* 2020 Aug;39(8):2368-88. <https://doi.org/10.1016/j.clnu.2019.11.024>[PubMed]1532-1983.
7. WHO. Obesity: preventing and managing the global epidemic. Report on a WHO Consultation on Obesity, Geneva, 3-5 June, 1997. WHO/NUT/NCD/98. 1. Technical Report Series Number 894. Geneva: World Health Organization, 2000.
8. Expert Consultation WH; WHO Expert Consultation. Appropriate body-mass index for Asian populations and its implications for policy and intervention strategies. [Erratum in: *Lancet.* 2004 Mar 13;363] [9412] [902. PMID: 14726171]. *Lancet.* 2004 Jan;363(9403):157-63. [https://doi.org/10.1016/S0140-6736\(03\)15268-3](https://doi.org/10.1016/S0140-6736(03)15268-3)[PubMed]1474-547X.
9. Shah NR, Braverman ER. Measuring adiposity in patients: the utility of body mass index (BMI), percent body fat, and leptin. *PLoS One.* 2012;7(4):e33308. <https://doi.org/10.1371/journal.pone.0033308>[PubMed]1932-6203.
10. Batsis JA, Mackenzie TA, Bartels SJ, et al. Diagnostic accuracy of body mass index to identify obesity in older adults: NHANES 1999-2004. *Int J Obes.* 2016 May;40(5):761-7. <https://doi.org/10.1038/sj.ijo.2015.243>[PubMed]1476-5497.
11. Examination Committee of Criteria for 'Obesity Disease' in Japan; Japan Society for the Study of Obesity. New criteria for 'obesity disease'. *Jpn Circ J.* 2002 Nov;66(11):987-92. <https://doi.org/10.1253/circj.66.9870047-1828>.
12. Lean ME, Han TS, Morrison CE. Waist circumference as a measure for indicating need for weight management. *BMJ.* 1995 Jul;311(6998):158-61. <https://doi.org/10.1136/bmj.311.6998.158>[PubMed]0959-8138.
13. National Institutes of Health. Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults – The Evidence Report. *Obes Res.* 1998 Sep;6 Suppl 2:51S-209S.[PubMed]1071-7323.
14. Misra A, Vikram NK, Gupta R, et al. Waist circumference cutoff points and action levels for Asian Indians for identification of abdominal obesity. *Int J Obes.* 2006 Jan;30(1):106-11. <https://doi.org/10.1038/sj.ijo.0803111>[PubMed]0307-0565.
15. Wang Z, Ma J, Si D. Optimal cut-off values and population means of waist circumference in different populations. *Nutr Res Rev.* 2010 Dec;23(2):191-9. <https://doi.org/10.1017/S0954422410000120>[PubMed]1475-2700.
16. Cruz-Jentoft AJ, Bahat G, Bauer J, et al. Sarcopenia: revised European consensus on definition and diagnosis. *Age and Ageing.* 2019; 48(1):16-31.
17. Dodds RM, Syddall HE, Cooper R, et al. Grip strength across the life course: normative data from twelve British studies. *PLoS One.* 2014 Dec;9(12):e113637. <https://doi.org/10.1371/journal.pone.0113637>[PubMed]1932-6203.
18. Bhasin S, Trivison TG, Manini TM, et al. Sarcopenia Definition: The Position Statements of the Sarcopenia Definition and Outcomes Consortium. *J Am Geriatr Soc.* 2020 Jul;68(7):1410-8. <https://doi.org/10.1111/jgs.16372>[PubMed]1532-5415.
19. Patel SM, Duchowny KA, Kiel DP, et al Sarcopenia Definition & Outcomes Consortium Defined Low Grip Strength in Two Cross-Sectional, Population-Based Cohorts. *J Am Geriatr Soc.* 2020 Jul;68(7):1438-44. <https://doi.org/10.1111/jgs.16419>[PubMed]1532-5415.
20. Lauretani F, Russo CR, Bandinelli S, Bartali B, Cavazzini C, Di Iorio A, et al Age-associated changes in skeletal muscles and their effect on mobility: an operational diagnosis of sarcopenia. *J Appl Physiol* (1985). 2003 Nov;95(5):1851-60. doi: <https://doi.org/10.1152/jappphysiol.00246.2003>
21. Studenski SA, Peters KW, Alley DE, et al The FNIH sarcopenia project: rationale, study description, conference recommendations, and final estimates. *J Gerontol A Biol Sci Med Sci.* 2014 May;69(5):547-58. <https://doi.org/10.1093/gerona/glu010>[PubMed]1758-535X.
22. Chen LK, Woo J, Assantachai P, et al Asian Working Group for Sarcopenia: 2019 Consensus Update on Sarcopenia Diagnosis and Treatment. *J Am Med Dir Assoc.* 2020 Mar;21(3):300-307.e2. <https://doi.org/10.1016/j.jamda.2019.12.012>[PubMed]1538-9375.
23. Auyeung TW, Arai H, Chen LK, Woo J. Letter to the editor: Normative data of handgrip strength in 26344 older adults – a pooled dataset from eight cohorts in Asia. *J Nutr Health Aging.* 2020;24(1):125-126. doi: <https://doi.org/10.1007/s12603-019-1287-6>.
24. Assantachai P, Muangpaisan W, Intalapaporn S, et al. Cut-off points of quadriceps strength, declines and relationships of sarcopenia-related variables among Thai community-dwelling older adults. *Geriatr Gerontol Int.* 2014 Feb;14 Suppl 1:61-8. <https://doi.org/10.1111/ggi.12207>[PubMed]1447-0594.
25. Martien S, Delecluse C, Boen F, et al Is knee extension strength a better predictor of functional performance than handgrip strength among older adults in three different settings? *Arch Gerontol Geriatr.* 2015 Mar-Apr;60(2):252-8. <https://doi.org/10.1016/j.archger.2014.11.010>[PubMed]1872-6976.
26. Cesari M, Kritchevsky SB, Newman AB, et al; Health, Aging and Body Composition Study. Added value of physical performance measures in predicting adverse health-related events: results from the Health, Aging And Body Composition Study. *J Am Geriatr Soc.* 2009 Feb;57(2):251-9. <https://doi.org/10.1111/j.1532-5415.2008.02126.x>[PubMed]1532-5415.
27. Gallagher D, Heymsfield SB, Heo M, Jebb SA, Murgatroyd PR, Sakamoto Y. Healthy percentage body fat ranges: an approach for developing guidelines based on body mass index. *Am J Clin Nutr.* 2000 Sep;72(3):694-701. <https://doi.org/10.1093/ajcn/72.3.694>[PubMed]0002-9165.
28. Baumgartner RN, Koehler KM, Gallagher D, et al Epidemiology of sarcopenia among the elderly in New Mexico. [Erratum in: *Am J Epidemiol* 1999 Jun 15;149] [12]. *Am J Epidemiol.* 1998 Apr;147(8):755-63. <https://doi.org/10.1093/oxfordjournals.aje.a009520>[PubMed]0002-9262.
29. Ishii S, Chang C, Tanaka T, et al The Association between Sarcopenic Obesity and Depressive Symptoms in Older Japanese Adults. *PLoS One.* 2016 Sep;11(9):e0162898. <https://doi.org/10.1371/journal.pone.0162898>[PubMed]1932-6203.
30. Bahat G, Kilic C, Topcu Y, et al. Fat percentage cutoff values to define obesity and prevalence of sarcopenic obesity in community-dwelling older adults in Turkey. *Aging Male.* 2020 Dec;23(5):477-2. <https://doi.org/10.1080/13685538.2018.1530208>[PubMed]1473-0790.
31. Pedrero-Chamizo R, Gómez-Cabello A, Meléndez A, et al. Higher levels of physical fitness are associated with a reduced risk of suffering sarcopenic obesity and better perceived health among the elderly: the EXERNET multi-center study. *J Nutr Health Aging.* 2015 Feb;19(2):211-7.
32. Kim TN, Yang SJ, Yoo HJ, et al. Prevalence of sarcopenia and sarcopenic obesity in Korean adults: the Korean sarcopenic obesity study. *Int J Obes.* 2009 Aug;33(8):885-92.
33. Janssen I, Heymsfield SB, Ross R. Low relative skeletal muscle mass (sarcopenia) in older persons is associated with functional impairment and physical disability. *J Am Geriatr Soc.* 2002 May;50(5):889-96. <https://doi.org/10.1046/j.1532-5415.2002.50216.x>[PubMed]0002-8614.
34. Lee J, Hong YP, Shin HJ, Lee W. Associations of Sarcopenia and Sarcopenic Obesity With Metabolic Syndrome Considering Both Muscle Mass and Muscle Strength. *J Prev Med Public Health.* 2016 Jan;49(1):35-44. <https://doi.org/10.3961/jpmph.15.055>[PubMed]2233-4521.
35. Zoico E, Di Francesco V, Guralnik JM, et al. Physical disability and muscular strength in relation to obesity and different body composition indexes in a sample of healthy elderly women. *Int J Obes.* 2004 Feb;28(2):234-41.
36. Lim S, Kim JH, Yoon JW, et al Sarcopenic obesity: prevalence and association with metabolic syndrome in the Korean Longitudinal Study on Health and Aging (KLoSHA). *Diabetes Care.* 2010 Jul;33(7):1652-4. <https://doi.org/10.2337/dc10-0107>[PubMed]1935-5548.
37. Baek SJ, Nam GE, Han KD, et al Sarcopenia and sarcopenic obesity and their association with dyslipidemia in Korean elderly men: the 2008-2010 Korea National Health and Nutrition Examination Survey. *J Endocrinol Invest.* 2014;37(3):247e60. <https://doi.org/10.1007/s40618-013-0011-3>.
38. Batsis JA, Barre LK, Mackenzie TA, et al. Variation in the prevalence of sarcopenia and sarcopenic obesity in older adults associated with different research definitions: dual-energy X-ray absorptiometry data from the National Health and Nutrition Examination Survey 1999e2004. *J Am Geriatr Soc.* 2013;61(6):974e80.
39. Chung JY, Kang HT, Lee DC, Lee HR, Lee YJ. Body composition and its association with cardiometabolic risk factors in the elderly: a focus on sarcopenic obesity. *Arch Gerontol Geriatr.* 2013 Jan-Feb;56(1):270-8.
40. Volkert D, Beck AM, Cederholm T, et al. ESPEN guideline on clinical nutrition and hydration in geriatrics. *Clin Nutr.* 2019;38(1):10-47.